

Use of Sterile Grass Carp to Control Aquatic Weeds

George W. Lewis
Professor, Aquaculture and Fisheries
Warnell School of Forest Resources
The University of Georgia

Revised May 19, 1998

Aquatic weeds are a serious problem for pond owners throughout Georgia. They restrict access to fishing areas, reduce fish harvest and decrease the usefulness, attractiveness and value of a pond. Herbicides, mechanical removal, water level changes, dyes, fertilization, proper pond construction, pond renovation and biological methods successfully control unwanted aquatic weed growth. The physical and chemical characteristics of the pond and the pond owner's objectives dictate which method is most appropriate. Pond owners are familiar with the traditional methods of aquatic weed control but not with the recently available option of biological control by using sterile triploid grass carp (Figure 1).




Figure 1. Nine-inch triploid grass carp stocked in a central Georgia pond with a heavy weed infestation grew to lengths of 29 inches and weights of almost 20 pounds in 16 months.

The grass carp (*Cteno pharyngodon idella*) occurs naturally in large rivers of the eastern USSR and China. It was introduced into the United States in 1963 by the United States Bureau of Sport Fisheries and Wildlife in cooperation with Auburn University. The feeding habits of the grass carp were well known and it was thought to have great potential as a biological weed control agent. However, there was concern that the grass carp could reproduce in the wild and become an environmental nuisance, destroying valuable areas such as wetlands, swamps and waterfowl feeding grounds.

Because of these environmental concerns, early research focused on developing sterile populations. Attempts included producing single-gender populations, creating sterile hybrids and removing gonads. Success was limited because these methods were seldom 100 percent effective and verification of sterility was difficult. In the early 1980s researchers and commercial producers began treating eggs with heat, cold or pressure to inhibit the second maturation division in the fertilized egg. This produced fish with abnormal chromosome numbers. The normal diploid grass carp has a chromosome number (2N) of 48, while the triploid grass carp has a chromosome number (3N) of 72. The extra chromosomes result in sterility. Unfortunately, not all treated eggs develop into fish with abnormal chromosome numbers. A technique using an electronic particle size analyzer was developed in the early 1980s which identifies carp as triploids or diploids. Georgia laws and regulations require each grass carp be verified and documented as a triploid.

Feeding Habits

Triploid and diploid grass carp appear to consume similar quantities of aquatic plants and to have similar feeding habits, preferring succulent young plants. Table 1 lists some common aquatic plants and rates them by grass carp feeding preferences. Grass carp will not control all types of aquatic weeds. Because of selective feeding habits, they can eliminate one plant species and make room for the expansion of others.

Table 1. Feeding preferences of grass carp on some common aquatic plants.

High	Moderate	Low
Musk-grass	Duckweeds	Eel grass
Naiads	Pondweeds	Watermeal
Hydrilla	Bladderwort	Cattail
American elodea	Fanwort	Milfoil
	Water pennywort	Parrot feather
	Coontail	Reeds
	Water primrose	Sedges
	Filamentous algae	Water hyacinth
		Alligator weed
		Spatterdock
		Yellow cowlily
		Maidencane
		Torpedo grass
		Watershield Waterlily

Stocking

If the grass carp is the preferred weed control option, stocking proper numbers is important. Stocking rates of five to over 200 fish per acre have been used depending on plant species, plant density and distribution, the size and age of the fish and the pond owner's objectives. There are computer models that determine the appropriate stocking density by considering additional factors such as the amount of human activity around the pond, the desired level of control and grass carp feeding preferences. The numbers recommended are designed to provide a 75 to 90 percent reduction in target plant species in three to four years. In most situations complete removal of aquatic vegetation is undesirable because the vegetation provides cover for small fishes and attachment surfaces for fish food organisms.

Grass carp stocking densities are based on the maximum expected weed coverage and the feeding preference rating of the weeds. Stock ten, 15 or 20 fish per acre depending on whether the target weed species is high, moderate or low on the feeding preference list, respectively. This stocking concept is best illustrated using a few examples:

Example 1: A 10 acre pond is examined in March and found to have five acres of naiads growing in it. However, three of the remaining five acres are shallow and the naiads are expected to spread to this area later in the growing season. Base the stocking rate on the maximum expected weed coverage (eight acres). Because naiads are high on the feeding preference list, stock a total of 80 fish (eight acres times 10 fish per acre).

Example 2: A 10 acre pond is examined in March and found to have five acres covered in watermeal. Because watermeal is a floating plant, pond depth does not matter. The maximum expected weed coverage would be the entire 10 acres. Stock two hundred fish (10 acres times 20 fish per acre).

Example 3: A 10 acre pond is examined in March and found to have a one acre infestation of water primrose. Because water primrose grows only in shallow water (less than two feet deep), base the stocking rate on the area of the pond less than two feet deep. If two acres of the pond are less than two feet deep, stock 30 fish (two acres times 15 fish per acre).

Grass carp could be stocked in weed-free ponds at low rates (five fish per acre) to prevent weeds from becoming established. However, the effectiveness of preventive stocking has not been determined. Generally, no fewer than 10 fish should ever be stocked, regardless of the pond size, because the loss of even a few fish could result in ineffective weed control.

The number of fish and the time required to achieve weed control can be reduced by using grass carp with other aquatic weed control options. For example, herbicides or mechanical removal can be used prior to fish introductions. If the established aquatic vegetation is removed, the tender new growth can be controlled by fewer fish.

Time of stocking affects the initial degree of weed control. Fishes are cold-blooded animals whose feeding rates and metabolism are influenced by water temperature. Grass carp feeding is greatest when the water temperature is between 70 and 80 degrees Fahrenheit and negligible when it is less than 50 degrees Fahrenheit. Mortality associated with handling stress is less likely when the water temperature is cooler; therefore, fish stocked in late winter or early spring are more likely to survive. They will not begin feeding heavily until late spring or early summer which is when most aquatic weeds begin growing in Georgia.

Because grass carp are attracted to currents, ponds with water flowing over spillways or through drains are not suitable without renovation. Cover horizontal drains with a fence or bars that allow free flow of water but prevent passage of grass carp. If barriers are placed over any drain structures, make sure they do not become clogged or blocked. Water could flow over emergency spillways and possibly wash out the spillway or dam.

Predatory fish, such as largemouth bass, eat grass carp. If used with existing fish populations, grass carp should be large enough to avoid being eaten by the average size predator. A largemouth bass 12 to 14 inches long can swallow a grass carp approximately nine inches long. Even if predation is not a problem, the pond owner should consider using larger carp if they are available because they tend to survive handling and stocking better. Grass carp stocked with existing fish populations should be at least 8-10 inches in length.

Grass carp do not reproduce in ponds and periodic restocking is required. It has been reported that the lifespan of the grass carp is between 10 and 15 years; however, triploid grass carp will provide effective vegetation control for 8-10 years.

Grass carp grow rapidly in ponds that have preferred plant species. Nine to 11 inch fish stocked in the early spring can reach lengths of 25 inches or more and weights of seven to 10 pounds by the end of the first year. If appropriate numbers of grass carp are stocked, they will eventually reduce the vegetation to the point that new plant growth is eaten as it becomes available. The grass carp will survive and remain healthy but will not increase in size. Once stocked, grass carp are difficult to remove from a pond. They are almost impossible to remove by seining or angling. The only options are draining the pond or using toxicants such as rotenone.

Permit Requirements

Only certain producers are authorized to sell grass carp in Georgia because they must confirm that each fish is a triploid. The Georgia Department of Natural Resources periodically examines shipments of grass carp to verify triploidy. If diploid fish are found, the person possessing or selling the fish is subject to serious legal action, including large fines and imprisonment, as well as having the stock

destroyed.

Using grass carp in Georgia does not require a permit from the Georgia Department of Natural Resources if the pond owner meets the following criteria:

1. Sterile triploid grass carp are purchased from sources authorized by the Department of Natural Resources. A list of currently approved dealers is available from county Extension and Georgia Department of Natural Resources offices.
2. The pond owner retains the bill of sale as proof of legal purchase.
3. The pond is privately owned, that is, a body of water which is clearly and entirely within the title of one owner.
4. Fish cannot travel upstream or downstream directly into a body of water not owned by the pond owner.

If unsure the above criteria can be met, consult a Department of Natural Resources fisheries biologist. Georgia Department of Natural Resources Fisheries offices are listed at the end of this publication.

Sources of Fish

Currently, there are several approved dealers for triploid grass carp in Georgia. Prices vary considerably depending on availability, fish size, season and dealer. Contact several of the approved sources for the best price.

Department of Natural Resources Wildlife Resources Division Fisheries Management Section

State Office

Chief of Fisheries
2070 US Highway 278, SE
Social Circle, GA 30025
770/918-6418

Regional Offices

Northwest Region I
312 North River St., NW
Calhoun, GA 30703-0519
706/624- 1161

Northern Region II
2150 Dawsonville Highway
Gainesville, GA 30501
770/535-5498

East Central Region III
2123 US Highway 278, SE
Social Circle, GA 30025
770/918-6418

West Central Region IV

1014 Martin Luther King Blvd
Fort Valley, GA 31030
478/825-6151

Southwest Region V
2024 Newton Road
Albany, GA 31701-3576
229/430-4256

South Central Region VI
108 Darling Avenue
Waycross, GA 31502-2089
912/285-6094

Coastal Region VII
22814 Highway 144
Richmond Hill, GA 31324
912/727/2112

District Offices

Bowens Mill Dist. Office
1773-A Bowens Mill Hwy.
Fitzgerald, GA 31760
229/426-5257

Metter Dist. Office
360 Cedar Street
Metter, GA 30439
912/684-6424

Thomson District Office
142 Bob Kirk Rd., NW
Thomson, GA 30824
706/595-5639